Head Injury Advice Sheet

Advice for parents and carers of children





How is your child?



If your child has any of the following during the next 48 hours:

- Vomits repeatedly i.e. more than twice (at least 10 minutes between each vomit)
- Becomes confused or unaware of their surroundings
- Loses consciousness, becomes drowsy or difficult to wake
- Has a convulsion or fit
- Develops difficulty speaking or understanding what you are saying
- Develops weakness in their arms and legs or starts losing their balance
- Develops problems with their eyesight
- Has clear fluid coming out of their nose or ears
- Does not wake for feeds or cries constantly and cannot be soothed

You need urgent help

Go to the nearest Hospital Emergency (A&E) Department or phone 999



If your child has any of the following during the next 48 hours:

- Develops a persistent headache that doesn't go away (despite painkillers such as paracetamol or ibuprofen)
- Develops a worsening headache

You need to contact a doctor or nurse today

Please ring your GP surgery or call NHS 111 - dial 111



If your child:

- Is alert and interacts with you
- Vomits, but only up to twice
- Experiences mild headaches, struggles to concentrate, lacks appetite or has problems sleeping

If you are very concerned about these symptoms or they go on for more than 2 months, make an appointment to see your GP.

Self Care
Continue providing
your child's care at
home. If you are still
concerned about your
child, call NHS 111 –
dial 111

How can I look after my child?

- Ensure that they have plenty of rest initially. A gradual return to normal activities/school is always recommended.
- Increase activities only as symptoms improve and at a manageable pace.
- It is best to avoid computer games, sporting activity and excessive exercise until all symptoms have improved.

Head Injury Advice Sheet

Advice for parents and carers of children





Concussion following a head injury

- Symptoms of concussion include mild headache, feeling sick (without vomiting), dizziness, bad temper, problems concentrating, difficulty remembering things, tiredness, lack of appetite or problems sleeping these can last for a few days, weeks or even months. Some symptoms resolve quickly whilst others may take a little longer.
- Concussion can happen after a mild head injury, even if they haven't been "knocked out".
- 9 out of 10 children with concussion recover fully, but some can experience long term effects, especially if they return to sporting activities too quickly. It is really important that your child has a gradual return to normal activities and that they are assessed by a doctor before beginning activities that may result in them having another head injury.
- If you are very concerned about these symptoms or they last longer than 2 months, you should seek medical advice from your doctor.

Advice about going back to nursery / school

- Don't allow your child to return to school until you feel that they have completely recovered.
- Try not to leave your child alone at home for the first 48 hours after a significant head injury.

Advice about returning to sport

- Repeated head injury during recovery from concussion can cause long term damage to a child's brain.
- Expect to stay off sport until at least 2 weeks after symptoms are fully recovered.
- Always discuss with your child's school and sports club to discuss a gradual return to full activity.

For further information:

Rugby: goo.gl/1fsBXz



Football: goo.gl/zAgbMx



For further support and advice about head injuries, contact:



■ Call their helpline on 0303 303 2248

Download factsheet from website: goo.gl/SPv4RJ

■ Visit their support page: goo.gl/r9PZth



